

Scaling Study of the Leptonic Decay Constants of Heavy-Light Mesons: A Consumers Report on Improvement Factors

S. Güsken^a, in collaboration with
C. Alexandrou^b, F. Jegerlehner^c, K. Schilling^a, G. Siegert^a and R. Sommer^d

^aPhysics Department, University of Wuppertal, D-42097 Wuppertal, Germany

^bDepartment of Natural Sciences, University of Cyprus, Nicosia, Cyprus

^cPSI, CH-5232 Villigen, Switzerland

^dDESY, Theory Division, D-22603 Hamburg, Germany

A high statistics calculation, performed at $\beta = 5.74, 6.00$ and 6.26 , enables us to study the variation of the leptonic decay constants f_P of heavy pseudoscalar mesons with the lattice spacing a . We observe only a weak a dependence when the standard $\sqrt{2\kappa}$ normalization is used for the quark fields, whereas application of the Kronfeld-Mackenzie normalization induces a stronger variation with a . Increasing the meson mass from 1.1GeV to 2.3GeV this situation becomes even more pronounced.

1. INTRODUCTION

The prediction of the leptonic decay constants of the D and the B meson within the framework of lattice QCD is a challenging but also very delicate problem, since in the region of heavy mesons, the inverse of the respective masses comes close to currently reachable lattice resolutions. Therefore large discretization effects may contaminate the results.

The question how to suppress these unphysical contributions has been tackled from various sides[1–3]. Using meanfield arguments, Kronfeld and Mackenzie suggested that the replacement of the standard $\sqrt{2\kappa}$ normalization of Wilson quark fields by $\sqrt{1 - 3\kappa/4\kappa_c}$ should significantly reduce the effects of finite lattice spacing.

The present study takes a rather empirical approach to the issue of finite a effects in the region of heavy mesons. We vary the lattice spacing in the currently accessible range and analyse the corresponding variation of f_P . Clearly a strong dependence of f_P on a would indicate the presence of large discretisation errors, whereas a weak dependence would be associated with smaller contaminations. In this sense our approach is perfectly suited to judge on the efficiency of different

quark field normalizations within a given a region.

We will finally perform an extrapolation of f_P to the continuum, assuming that its functional dependence on a is linear in the leading part, as suggested by the data.

2. PREPARATION

In order to visualize unambiguously the a dependence of f_P , we have to take care that the finite a effects are not hidden in the statistical noise or distorted by incomplete groundstate projection of the meson propagator and effects due to the finite size of the lattice. Therefore we have done our calculation with high statistics, keeping the errors of the raw data below 5%. We have varied the lattice size from about 0.7 fm to 2 fm and have smeared the quark fields with the well established [4] Gauss like Wuppertal wavefunction ($n = 100, \alpha = 4$). In table 1 we display the lattice parameters together with the lattice spacing, taken from the stringtension σ [5]. The influence of the finite lattice extension on f_P was checked by comparing the results at different lattice sizes and fixed lattice constant. We find that finite size effects are small once the lattice extension becomes as large as 1.4 fm .

Table 1
Lattice parameters

$\beta = 5.74, a_\sigma^{-1} = 1.118(9)$		$\beta = 6.00, a_\sigma^{-1} = 1.876(19)$		$\beta = 6.26, a_\sigma^{-1} = 2.775(18)$	
N_S, N_T	<i>no.conf.s.</i>	N_S, N_T	<i>no.conf.s.</i>	N_S, N_T	<i>no.conf.s.</i>
4,24	404	6,36	227		
6,24	131			12,48	103
8,24	175	12,36	204	18,48	76
10,24	213	18,36	9		

3. RESULTS

3.1. Finite a effects

In fig.1 we show the leptonic decay constant f_P as a function of a both in the $\sqrt{2\kappa}$ normalization (open symbols) and in the Kronfeld-Mackenzie normalization (closed symbols). The light quark




Figure 1. f_P as a function of a . Points connected only by dashed lines were not used in the extrapolation.

mass has been extrapolated to the chiral limit and we have interpolated between the results at adjacent heavy quark masses (c.f. table 2) in order to keep the meson mass (in GeV) fixed when the a dependence of $f_P(M_P, a)$ is investigated. Due to their small statistical errors, we have used stringtension measurements [5] to relate our data

to a physical scale. The renormalization factor Z_A was taken from perturbation theory [6]¹, with an effective coupling $\tilde{g}^2 = 3g_0^2 / <tr P_{\mu\nu}>$, recommended in ref.[3] ($P_{\mu\nu} \equiv 1 \times 1$ Wilson loop).

It goes without saying that the a dependence of f_P must be different in the two normalizations. Very surprisingly, however, fig.1 shows clearly that - in contrast to the $\sqrt{2\kappa}$ normalized results - the variation with a becomes stronger and stronger with increasing meson mass when the KroMac normalization is used. This means that - at least in the displayed a and M_P range - the KroMac normalization does a bad job: Instead of suppressing finite a effects it enhances them.

In order to connect our results to the (physical) continuum, we followed the behavior suggested by the data in both normalizations and extrapolated linearly² to $a = 0$. As can be seen from fig.1, we obtain nice agreement of the results, although the KroMac normalization has induced considerably larger errors.

3.2. Heavy mass extrapolation

The most 'natural' scale for f_P is f_π , since the uncertainty originating from the renormalization constant Z_A cancels out in this case. Lattice measurements of f_π are generally affected with large statistical errors and therefore we have decided to convert our results to this scale only after having performed the $a \rightarrow 0$ extrapolation of f_P . To achieve this we have decoupled the extrapolations according to $\frac{f_P}{f_\pi}(a \rightarrow 0) = \frac{f_P/\sqrt{\sigma}(a \rightarrow 0)}{f_\pi/\sqrt{\sigma}(a \rightarrow 0)}$.

¹ $Z_A = 1 - 0.1333g^2$ for standard normalization and $Z_A = 1 - 0.0248g^2$ in the case of KroMac normalization.

² Since the a dependence cannot be exactly linear for both normalizations at the same time, we have excluded those points from the fit where $\sqrt{1 - 3\frac{\kappa}{4\kappa_c}}/\sqrt{2\kappa} > 1.6$.

Table 2

Decay constant and meson mass in lattice units. The light quark has been extrapolated to κ_c .

$\beta = 5.74$			$\beta = 6.00$			$\beta = 6.26$		
κ_h	f_P/Z_A	M_P	κ_h	f_P/Z_A	M_P	κ_h	f_P/Z_A	M_P
0.06	0.1197(102)	2.502(13)	0.10	0.0873(17)	1.498(10)	0.09	0.0437(32)	1.579(19)
0.09	0.1629(52)	1.871(13)	0.115	0.0983(18)	1.197(7)	0.10	0.0486(31)	1.375(14)
0.125	0.1890(33)	1.205(7)	0.125	0.1038(18)	0.995(5)	0.120	0.0609(33)	0.965(10)
0.140	0.1907(38)	0.904(6)	0.135	0.1085(19)	0.780(7)	0.135	0.0689(29)	0.636(6)
0.150	0.1829(53)	0.684(5)	0.145	0.1032(31)	0.551(2)	0.145	0.0711(24)	0.382(4)
						0.1492	0.0580(50)	0.245(4)

Although the $O(\tilde{g}^4)$ -uncertainty in Z_A does not cancel out exactly if one first extrapolates and then takes the ratio, its effect should be roughly the same in numerator and denominator. To obtain the denominator of this ratio we used both our own data and the results quoted in refs. [7,8,2,9]. Since the a dependence of $f_\pi/\sqrt{\sigma}$ is weak, a linear extrapolation to $a = 0$ is well justified and leads to $\frac{f_\pi}{\sqrt{\sigma}}(a = 0) = 0.269(12)$.

In figure 2 we display our final results³ at $a = 0$ in the form⁴ $\hat{f}_P(1/M_P)$, together with our static value from ref. [4]. The new data appears to de-

points carry error bars of order 25% and therefore do not exclude a stronger variation in M_P . Given this situation we draw an error band that links the conventional results with the static point. The M_P dependence of the error band was chosen according to the ansatz $\hat{f}_P = c_0 + \frac{c_1}{M_P} + \frac{c_2}{M_P^2}$. At the location of the B and D meson the error band corresponds to the bounds $155MeV \leq f_B \leq 242MeV$, $150MeV \leq f_D \leq 200MeV$. It is evident from figure 2 that these bounds are strongly affected by the size and uncertainty of f_{stat} . More work is necessary to obtain an accurate prediction for f_B .




Figure 2. $\hat{f}$ as a function of $1/M_P$.

pend only weakly on M_P . Because of the various extra- and interpolations however, the data

³The extrapolation has been performed on the data in the $\sqrt{2\kappa}$ norm since it involves smaller statistical errors than using the KroMac norm.

⁴ $\hat{f} = f_P \sqrt{M_P} \times (\frac{\alpha_s(M_P)}{\alpha_s(M_B)})^{6/33}$.

REFERENCES

1. B. Sheikholeslami and R. Wohlert, Nucl. Phys. B259 (1985) 572.
2. C. Bernard et al., preprint UW/PT-93-06, Wash. U HEP/93-30, BNL-49068, 1993
3. G.P. Lepage, Nucl. Phys. B (Proc.Suppl.) 26 (1992) 45; A.S. Kronfeld, Nucl. Phys. B (Proc. Suppl.) 30 (1993) 445; P.B. Mackenzie, Nucl. Phys. B (Proc. Suppl.) 30 (1993) 35.
4. C. Alexandrou et al., prepr., PSI-92-27, 1992.
5. G.S. Bali and K. Schilling, Phys. Rev. D46 (1992) 2636; Wuppertal University preprint WUB-92-29, to appear in Phys. Rev. D.
6. R. Groot et al., Nucl. Phys. B237 (1984) 111.
7. F. Butler et al., Phys. Rev. Lett. 70 (1993) 2849; D. Weingarten, private communication.
8. The APE Collaboration, Rome preprint 93/928.
9. UKQCD Collaboration, Edinburgh Preprint 93/524.